

# Distribution and abundance of spiny lobster in the KWNWR and GWHNWR: Impacts of an intense recreational fishery

D. Eggleston, E. Johnson, C. Dahlgren,  
T. Kellison, D. Nadeau

**NC STATE UNIVERSITY**

# Objectives

- What is the magnitude of the impact of 2 d mini-season?
- What is the relationship between lobster density and fishing effort?


# KWNWR (2000)

## Three locations






- Lakes
- Marquesas
- Closed and Fished areas

## Diver Surveys


- Band transects
- Timed surveys




# Distribution and abundance by habitat




# Lobster density and sponges




# Impact of mini-season - KWNWR



# Impact of mini-season (Closed vs. Open)




# Effect of fishing effort







# GWHNWR (2002)





# Lobster density ( $> 76.2$ mm) before and after mini- season

- Patch Reefs ~ 95% decline
- Patch Heads ~ 79% decline
- 55% at Looe Key (Blonder et al. 1988)
- 58% in the Tortugas (Davis et al. 1977)



# Mini-season Effort



# Conclusions 1999

- In 1999, ~ 80% removal across all locations including 3 closed areas
- Lobster density was highest in Channels, Hardbottom, and Patch Reefs
- Lobster density was positively correlated to sponge area and volume

# Conclusions 2002

- In 2002, ~ 90% removal across all locations
- Fishing effort was 10 times greater on patch heads than patch reefs
- Fishing effort during mini-season was ~900 times greater than regular season
- Patterns in 2004 were similar, surveys will continue in 2005